
## 2015 Recommended Immunizations for Adults: By Age



There are several flu vaccines available. Talk to your healthcare professional about which flu vaccines is right for you.

pregnant, you should get a Tdap vaccine during the 3rd trimester of every pregnancy to help protect your babies from pertussis (whooping

cough).

You should get zoster vaccine even if you've had shingles before. There are two different types of pneumococcal vaccine: PCV13 (conjugate) and PPSV23 (polysaccharide). Talk with your healthcare professional to find out if one or both pneumococcal vaccines are recommended for you.

Your healthcare professional will let you know how many doses you need.

If you were born

in 1957 or after,

about how many doses you may need.

and don't have a

There are two HPV vaccines but only one HPV vaccine (Gardasil®)

record of being
vaccinated or
having had
measles, mumps
and rubella,
talk to your
healthcare
professional

If you are a male 22 through 26
years old and have sex with men
you should complete the HPV
vaccine series if you have not
already done so.

should be given to men.

Your healthcare professional will let you know how many doses you need.

**Recommended For You:** This vaccine is recommended for you *unless* your healthcare professional tells you that you cannot safely receive it or that you do not need it.



**May Be Recommended For You:** This vaccine is recommended for you if you have certain risk factors due to your health, job, or lifestyle that are not listed here. Talk to your healthcare professional to see if you need this vaccine.

If you are traveling outside the United States, you may need additional vaccines.

Ask your healthcare professional about which vaccines you may need at least 6 weeks prior to your travel.

For more information, call 1-800-CDC-INFO (1-800-232-4636) or visit www.cdc.gov/vaccines



U.S. Department of Health and Human Services Centers for Disease Control and Prevention

## 2015 Recommended Immunizations for Adults: By Health Condition

If you have this health condition. talk to your healthcare professional about these vaccines Flu Td/Tdap **Shingles Pneumococcal** Meningococcal MMR **HPV** Chickenpox Hepatitis A Hepatitis B Hib Influenza Tetanus, Zoster Measles, Human papillomavirus Varicella Haemophilus influenzae diphtheria, mumps, pertussis PCV13 PPSV23 rubella for women for men type b \*see below 1 - 2 doses Pregnancy 3 doses post-HSCT\* **SHOULD SHOULD SHOULD** Weakened Immune through **NOT GET** NOT GET **NOT GET** 3 doses age 26 years System recipients only VACCINE VACCINE VACCINE HIV: CD4 count less more than 200 doses HIV: CD4 count 200 1 dose 1 dose 1 or 3 of Tdap or greater 2 doses 3 doses doses 3 doses followed Flu Kidney disease or by Td booster 3 doses through age 21 vaccine through doses poor kidney function every age 26 every 10 years year 2 doses years years Asplenia (if you do not 1 dose 1 or for those 1 or 2 1 or 3 have a spleen or if it more doses doses 60 years doses does not work well) 3 doses or older Heart disease Chronic lung disease 1 or 3 Chronic alcoholism doses 1 dose Diabetes (Type 1 or Type 2) 3 doses doses Chronic Liver Disease 2 doses

**More Information:** 

There are several flu vaccines available. Talk to your healthcare professional about which flu vaccines is right for you.

\* If you are pregnant, you should get a Tdap vaccine during the 3rd trimester of every pregnancy to help protect your babies from pertussis (whooping

cough).

You should

get zoster
vaccine even
if you've
had shingles
before.

There are two different types of pneumococcal vaccine: PCV13 (conjugate) and PPSV23 (polysaccharide). Talk with your healthcare professional to find out if one or both pneumococcal vaccines are recommended for you.

Your healthcare professional will let you know how many doses you need.

If you were born

in 1957 or after.

about how many doses you

may need.

and don't have a

There are two HPV vaccines but only one HPV vaccine (Gardasil®)

should be given to men.

Recommended for you if you did not get it when you were a child.


record of being
vaccinated or
having had
measles, mumps
and rubella,
talk to your
healthcare
professional

If you are a male 22 through 26
years old and have sex with men
you should complete the HPV
vaccine series if you have not
already done so.

Your healthcare professional will let you know how many doses you need.

\*Hematopoietic stem cell transplant

**Recommended For You:** This vaccine is recommended for you *unless* your healthcare professional tells you that you cannot safely receive it or that you do not need it.



YOU SHOULD NOT GET THIS VACCINE

If you are traveling outside the United States, you may need additional vaccines.

Ask your healthcare professional about which vaccines you may need at least 6 weeks prior to your travel.

For more information, call 1-800-CDC-INFO (1-800-232-4636) or visit www.cdc.gov/vaccines



U.S. Department of Health and Human Services Centers for Disease Control and Prevention